

LIFE PINNARCA

LIFE NAT/ES/001265



DELIVERABLE DC1.4

TITLE: DATA ON POSSIBLE LOCATIONS OF SOURCE
POPULATIONS OF LARVAE FOR THE PROJECT

ACTION C1: INSTALLATION OF LARVAL COLLECTORS

31/10/2024



RESPONSIBLE

PARTICIPANT 1
PARTICIPANT 2
PARTICIPANT 3
PARTICIPANT 4
PARTICIPANT 5
PARTICIPANT 6

CSIC

IMEDMAR-UCV
IRTA
IOPR
UA-CIMAR
UAGEAN
UNINA

To be cited as:

A. Feria-Rodríguez, D. March, B. Mourre, I.E. Hendriks, M. Vázquez-Luis. Sink-source connectivity for restocking of *Pinna nobilis* in the western Mediterranean Sea. *Marine Environmental Research*, Volume 197, 2024, 106428, <https://doi.org/10.1016/j.marenvres.2024.106428>.

Deliverable:

A. Feria-Rodríguez, D. March, B. Mourre, I.E. Hendriks, M. Vázquez-Luis. Sink-source connectivity for restocking of *Pinna nobilis* in the western Mediterranean. Report of possible locations of source populations of *Pinna nobilis* larvae. Within the Action C1: Installation of larval collectors. LIFE Pinnarca NAT/ES/001265 “Protection and restoration of *Pinna nobilis* populations as a response to the catastrophic pandemic started in 2016”, 31 pp.

This report is confidential, and the reproduction or dissemination of the content and data of this document to non-contracting third parties is prohibited, without the express authorization of partners, unless otherwise agreed by the parties granting the agreement when it is the object.



INDEX

Abstract	1
INTRODUCTION	2
MATERIALS & METHODS	6
Study area and data collection	6
Larval dispersal model	7
Connectivity analysis	8
RESULTS	10
DISCUSSION	14
References	19
Appendix 1	28

Abstract

The critically endangered endemic bivalve *Pinna nobilis* from the Mediterranean Sea suffered a sudden population decline after a mass mortality event in early autumn 2016. Conservation efforts aimed at preventing extinction included safeguarding resistant individuals and implementing a breeding plan to contribute to the repopulation of the species. This study utilized a model combining Lagrangian dispersion and connectivity analyses to pinpoint optimal restocking sites in the Western Mediterranean. Our approach allowed to identify locations capable of sustaining and generating larvae for broader repopulation in key areas of the Western Mediterranean Sea prior to the mass mortality event. Six important repopulation locations from Murcia, Valencia and Balearic Islands were selected for reintroduction efforts. The results obtained in this study show how the network could be self-sufficient and able to self-replenish itself of recruits. Overall, our work can be used to direct the reintroduction of resistant animals in the Western Mediterranean Sea.

